



Article Side

About diabetes | Symptoms of diabetes in women | Treatment of diabetes | Diabetes symptoms in women. by [Jacob Oram](#)

Article published on May 29th 2012 | [Health](#)

Other kinds of mesothelioma cancer consist of the more typical dangerous pleural mesothelioma cancer, occurring in the upper body walls tissue layer surrounding the respiratory system, and dangerous pericardial mesothelioma cancer, which occurs in the pericardial lining of the heart. Peritoneal mesothelioma cancer is the second most rare form of the condition and accounts for approximately 20% of all mesothelioma cancer situations each season.

What Causes Peritoneal Mesothelioma?

Peritoneal mesothelioma cancer is known only to be due to experience asbestos components fiber, a minute natural fiber that was used heavily in industry. Asbestos is said to reach the abdomen walls by one of two techniques. The first is through consumed asbestos components fiber components which are processed through digestive function and become filed in the peritoneum. The other method is through the respiratory system and lymph nodes, by consumed asbestos components fiber components. Malignant mesothelioma cancer of the pleura is also known to spread directly into the abdomen hole if its spread is not stunted.

Many of those who have worked with asbestos components fiber components have developed peritoneal mesothelioma cancer as a result of experience friable asbestos components fiber components. Asbestos was used in myriad industrial components and construction products. Visibility to asbestos components fiber was typical in metal vegetation, steel generators, power vegetation, and naval shipyards.

Peritoneal mesothelioma cancer is a relatively rare condition, with only 200-500 conclusions each season in the United States being recorded. Danger however, is continual in those who have been revealed to asbestos components fiber. Even those revealed many decades ago are possibly vulnerable for the progression of dangerous peritoneal mesothelioma cancer. Latency between experience asbestos components fiber and the progression of peritoneal mesothelioma cancer can be anywhere from 20-50 decades. Misdiagnosis is also typical because peritoneal mesothelioma cancer closely simulate minor stomach/abdominal discomfort and hernia. Nevertheless, those with clear experience asbestos components fiber history should be aware of the condition and speak with their doctor or cancer specialist if they believe they may be experiencing signs related to mesothelioma cancer.

How is Peritoneal Mesothelioma Treated?

Article Source:

<http://www.articleside.com/health-articles/about-diabetes-symptoms-of-diabetes-in-women-treatment-of-diabetes-diabetes-symptoms-in-women.htm> - [Article Side](#)

[Jacob Oram](#) - About Author:

those with clear experience asbestos components fiber history should be aware of the condition and speak with their doctor or cancer specialist if they believe they may be experiencing signs related to mesothelioma cancer.

How is Peritoneal Mesothelioma Treated?

Article Keywords:

low blood sugar, glucose, insulin levels,

You can find more [free articles](#) on [Article Side](#). Sign up today and share your knowledge to the community! It is completely FREE!