



Article published on May 1st 2012 | [Health](#)

A longstanding issue among certain females is breast asymmetry-it causes a woman to have mismatched breasts, or in other words, one of the breasts is notably smaller or larger than the other. Women of all ages can experience this.

Usually, breasts manifest themselves during the course of a girl's puberty, developing in time. However, this is not the case for one-fourth of the female population worldwide who suffer from breast asymmetry. Studies suggest that the typical weight of a breast should register at 16.9 ounces, and women having the condition would be self-conscious because of their light breasts.

Additionally, an uneven breast could be an underlying indication of breast cancer. According to the latest journal shared on Breast Cancer Research, women who possess asymmetrical breasts have a 50 percent risk of acquiring cancer of the breast. That is why it's vital for females with this issue to have doctors look at their asymmetrical breasts to find out if there are complications they need to be cautious about. Experts, however, noted that it's common for women to have slightly different breast shapes or sizes; what is worrying is similarly-sized breasts gradually becoming asymmetrical.

Genetics

Physicians report that genetics or haphazard growth patterns in the body are a contributing component why some women have breast asymmetry. The human breast develops due to the female hormone estrogen, growing two years after the girl experiences her very first menstrual period. Within the next 2 to 4 years, the breast continues expanding right until it attains maturity. It's not the truth for women who have irregular breast growths.

Cancer of the Breast

As stated by health news accounts, an irregular breast can reflect imminent health conditions, such as breast cancer. Doctors frequently propose breast augmentation Los Angeles health offices provide to counter the consequences of breast asymmetry, but self-breast examinations and regular mammogram screening should be performed as necessary.

Pregnancy and Sagging

There are tons of hormonal changes taking place in a pregnant woman's body. The breasts become more sensitive throughout the pregnancy phase. She might discover that her left breast is bigger than the other as a result of the production of milk, or lactation. On the other hand, ptosis or sagging happens when a woman begins to get older over time. Older women who went through breast augmentation procedures could also find their breasts sagging. This can be resolved with breast implant revision done on both breasts to improve their general physical appearance. Visit wdcyber.com/nbreast3.htm for more info about the prevalent reasons for breast asymmetry.

Article Source:

<http://www.articleside.com/health-articles/basic-facts-about-asymmetrical-breasts.htm> - [Article Side](#)

[Terry Bayer](#) - About Author:

For more details, search a [breast implant revision](#), a [breast augmentation Los Angeles](#) and a [breast](#)

[asymmetry](#) in Google for more related information.

Article Keywords:

breast asymmetry, breast augmentation los angeles, breast implant revision

You can find more [free articles](#) on [Article Side](#). Sign up today and share your knowledge to the community! It is completely FREE!